

NGM282 Promotes HDL Biogenesis and Transhepatic Cholesterol Efflux to Prevent Atherosclerosis in Mice

Lei Ling, Mei Zhou, R. Marc Learned, Hui Tian and Alex M DePaoli
NGM Biopharmaceuticals, South San Francisco, CA, United States




BACKGROUND AND AIMS

- FGF19, an endocrine hormone produced in the gut, acts in the liver to control bile acid synthesis¹⁻²
- NGM282 is an engineered, non-tumorigenic analogue of human FGF19³⁻⁴
- In phase 2 clinical trials in patients with non-alcoholic steatohepatitis, administration of NGM282 resulted in rapid and profound reductions in liver fat content, liver injury, inflammation and fibrosis⁵⁻⁶
- However, NGM282 increased cholesterol levels, and the molecular mechanisms that integrate the FGF19 signaling with cholesterol metabolic pathways are incompletely understood
- Here we investigate these mechanisms using a combination of pharmacological, bioinformatics, metabolomics and biochemical approaches

METHODS


- *db/db* mice (Jax#000642) received an intravenous injection of 1×10^{11} vector genome adeno-associated virus (AAV) carrying FGF19, NGM282 or green fluorescent protein (control). 2 weeks later, serum levels of cholesterol, HDL-C and LDL-C were measured. Livers were collected for transcriptome profiling, QPCR, metabolomics and histological analysis
- Hepatocyte-specific ABCA1-deficient mice were obtained by injecting *Abca1^{fl/fl}Abcg1^{fl/fl}* mice with AAV-TBG-Cre, which drives Cre recombinase expression under TBG promoter, allowing hepatocyte-specific expression
- FGFR4-deficient mice or wild-type mice received an intravenous injection of 3×10^{11} vector genome AAV carrying FGF19 or green fluorescent protein (control)
- ApoE-deficient mice were placed on a high-fat, high-cholesterol Western diet (Teklad TD88137) immediately following AAV injection, and this diet was continued ad libitum throughout the study. Mice were euthanized 18 weeks after AAV administration for analysis
- En face analyses of the ApoE-deficient mice were conducted at Wake Forest School of Medicine Metabolic Core (Winston-Salem, NC). Aortic lesion area was quantified using Image J software and expressed as percent stained area relative to total aortic area. All quantifications were carried out by an observer blinded to the sample identity
- Concentrations of total cholesterol, HDL-C and LDL-C were measured by enzymatic methods on an automated analyzer (COBAS INTEGRA 400 Plus Clinical Analyzer, Roche Diagnostics)



RESULTS




NGM282 Increases Serum Cholesterol in *db/db* Mice

- Administration of FGF19 or NGM282 resulted in elevations in total cholesterol, HDL-C and LDL-C in 2 weeks in *db/db* mice



NGM282 Selectively Activates Hepatic LXR Signaling in *db/db* Mice


- Transcriptome profiling and Ingenuity pathway analysis revealed that FGF19 and NGM282 activate the LXR/RXR pathway
- Metabolomics analysis uncovered an increase in intrahepatic hydroxysterols, natural ligands for LXR⁷
- However, FGF19 and NGM282 selectively modulate LXR signaling and upregulate genes in transhepatic cholesterol efflux without causing steatosis




(Oil droplets are shown as black vacuoles in livers stained with osmium tetroxide)

Deficiency in Hepatocellular ABCA1 Abolishes NGM282-Associated Cholesterol Change

- FGF19 and NGM282 promote HDL biogenesis through the induction of *Abca1* and *Apoa1*, both are LXR targets




- NGM282-associated cholesterol increases were blunted in mice deficient in ABCA1




NGM282-Associated Cholesterol Change is Dependent on FGFR4

- FGF19-associated cholesterol increases were abolished in mice deficient in FGFR4





- A constitutively active MEK1, but not a constitutively active STAT3, mimics the effect of FGF19 on cholesterol change



NGM282 Protects Against Atherosclerosis in ApoE-deficient Mice


- In dyslipidemic ApoE-deficient mice fed a Western diet, treatment with NGM282 significantly reduced atherosclerotic lesion area in aortas




LXR Agonist Increases Serum LDL-C in Humans

- Activation of LXR signaling is well known to promote reverse transport of cholesterol from peripheral tissues to the liver⁸
- LXR activation induces the expression of genes involved in cholesterol efflux, facilitates cholesterol esterification by promoting fatty acid synthesis, and exhibits anti-inflammatory effects through inhibition of TLR signaling⁸
- The development of LXR agonists as anti-atherogenic agents has been hindered by hepatic steatosis that accompanies activation of LXR in the liver⁹. Furthermore, first-in-human testing of LXR agonists revealed LDL-C elevations¹⁰
- Similarly, administration of NGM282 in patients with nonalcoholic steatohepatitis resulted in elevations in serum LDL-C levels, but reductions in liver fat content⁵

LXR Agonist BMS-852927 Increases LDL-C in Humans



Summary



CONCLUSION

- The endocrine hormone FGF19 and its analogue NGM282 have a hitherto unsuspected intrinsic role in promoting transhepatic cholesterol efflux and HDL biogenesis through selectively activating LXR signaling, while ameliorating steatosis and atherosclerosis
- The selective modulation of LXR by NGM282 may provide additional understanding of the observed cholesterol increases in animals and humans, as well as the cardiovascular benefit in animals

References:

1. Kliewer et al, *Dig Dis* 2015; 33:327-331
2. Degirolamo et al, *Nat Rev Drug Discov* 2016; 15:51-69
3. Zhou et al., *Cancer Res* 2014; 74:3306-3316
4. Luo et al., *Sci Transl Med* 2014; 6:247ra100
5. Harrison et al., *Lancet* 2018; 391:1174-1185
6. Harrison et al., *Hepatology* 2019; doi: 10.1002/hep.30590
7. Janowski et al., *Nature* 1996; 383:728-731
8. Tontonoz et al., *Mol Endocrinol* 2003; 17:985-93
9. Schultz et al., *Genes & development* 2000; 14:2831-2838
10. Kirchgessner et al., *Cell Metab* 2016; 24:223-33

This study was funded by NGM Biopharmaceuticals
Author disclosures on file at EASL 2019



DOI: 10.2455/psa.ea.LC2019.2019

General hepatology
Lei Ling



LBP-019