

Living Donor Kidney Transplantation(LDKT) from HBsAg (+) Donor to HBSAg (-) Recipient

Jin Min Kong¹, Sung Hyun Son¹, Hyukyong Kwon¹, Joon Heun Jeong², Eun Joo Hwang³, Won Young Choi³, Chul Soo Yoon², Chang Yell Lee², Dong Ryul Lee⁴, Jungmyung Ahn⁴

¹Nephrology, Hanseo Hospital, Busan, Korea; ²Surgery, Hanseo Hospital, Busan, Korea;

³Pathology, Hanseo Hospital, Busan, Korea; ⁴Nephrology, Marynoll Hospital, Busan, Korea.

Introduction

- HBsAg positivity is currently regarded as a contraindication of kidney donation to HBsAg negative recipients due to the risk of HBV transmission.
- Donor organ shortage is serious on this planet. The use of the HBsAg(+) donor organ may help to increase the opportunity of transplantation.
- We developed a protocol that enables LDKT from a HBsAg (+) donor to a HBsAg(-) recipient.

Protocol

- 1) Hepatitis B vaccine in recipients without protective titer (≥ 10 mIU/ml) of anti-HBs
- 2) Entecavir in donors with detectable HBV DNA in serum to reduce the viral load for 1-2 months before transplantation
- 3) HBIG in recipients on transplant day
- 4) Entecavir prophylaxis in recipients for 2-3 post-transplant months
- 5) Monitoring of HBV transmission in recipient ; HBsAg/anti-HBs titer, AST/ALT, and HBV Real Time-PCR on 1 day, 1 & 6 months after KT

Results

- Six LDKT from a HBsAg(+) donor to a HBsAg(-) recipient in 5 patients
- In one case, the recipient had a low titer of anti-HBs (7mIU/ml), which was raised by hepatitis B vaccine to 15mIU/ml before transplantation.
- All recipients had undetectable HBV DNA after LDKT and maintained HBsAg(-)/anti-HBs(+) status throughout the median follow up of 21(8-40) months

Conclusions

- Kidneys from HBsAg (+) living donors can be safely transplanted to HBsAg (-) recipients.

Table 1. Donors

Case #	1	2	3	4	5	6
HBsAg/Ab	+/-	+/-	+/-	+/-	+/-	+/-
HBeAg/Ab	-/+	-/+	-/+	-/+	NA*	-/+
Anti HBc	+	NA	+	+	+	+
HBV DNA	1.2×10^4	3.1×10^4	UD**	3.0×10^3	1.9×10^3	3.6×10^2
Use of Entecavir	Yes	Yes	No	Yes	Yes	Yes
HBV DNA after entecavir	UD	UD		UD	UD	UD

Table 2. Recipients

Case #	1	2	3	4	5	6	
HBsAg/Ab	Before KT	-/+	-/+	-/+	-/+	-/+	-/+
	Last measured after KT	-/+	-/+	-/+	-/+	-/+	-/+
HB Vaccine before KT	No	No	No	No	Yes	No	
Anti-HBs titer before KT	2,470	919	143	161	7 → 15 [†]	173	
HBV DNA, last measured	UD	UD	UD	UD	UD	UD	
Follow up months	40	33	21	20	20	8	

* NA; not available, ** UD; undetectable (< 20 IU/ml by real-time PCR), [†] Before and after vaccination

