

Urinary Kidney injury molecule-1 can predict acute tubular necrosis and delayed graft function in living donor renal allograft recipients


Narayan Prasad¹, Brijesh Yadav¹, Vikas Agarwal², Akhilesh Jaiswal¹, Vinita Agrawal³, R K Sharma¹
¹Department of Nephrology, ²Clinical Immunology, and ³ Pathology, Sanjay Gandhi Postgraduate Institute of Medical Sciences, Lucknow, India

Introduction

- Successful graft function for a long time periods depends on early and delayed graft function
- Delayed graft function (DGF)
 - >A complication of early impaired graft function
 - ➤ Defined by <70% decline in serum creatinine level within 48hours of transplant,
 - Need of dialysis within 7 days after transplant
- Prevalence of DGF
 - In live related transplant is (5-10%) and
 - In cadaveric related transplant (3-40%).

• Cause for DGF

- ✓ Acute tubular necrosis
- Consequence of "pH paradox"
- Free radical and reactive oxygen species mediated damage to renal tubules
- •Mechanism of acute tubular necrosis(Fig.1)



• Kidney injury molecule-1

- Single pass transmembrane glycoprotein
- Immunoglobulin like extracellular mucine domain
- The expression of KIM-1 upregulated after proximal tubules damage for its repairment
- Matrix metalloproteinase cleaves ectodomaine and released in urine
- Till now the available gold standard techniques is
- > 1st Serum creatinine
 - Expressed very late until a significant damage to kidney
 - ✓ Depends on muscle mass of patient
 - ✓ Depends on patient age, gender and ethnicity
- ▶ 2nd Biopsy of transplanted kidney
 - ✓ It is a invasive and
 - ✓ Increases the complication & jeopardized graft

Objectives

- 1. To determine the uKIM-1 level in urine sample of renal transplant recipient patients
- 2. To determine the specificity, Sensitivity, and predictive cutoff value of uKIM-1 for predicting DGF and ATN in live related renal allograft recipient
- 3. To determine the correlation between uKIM-1 and confounding factors of patients
- 4. To determine the suitability of uKIM-1 as diagnostic markers for DGF.

Materials and Methods

Patients recruitment

Live related renal transplant patient (n=56)

◆ Immediate graft function=47 ✓>70% Decline in serum creatinine within 48 hrs of transplant

transplant

✓ Patient have immediate post transplant urination >7litres

◆Delayed graft function=9 ✓ Biopsy proven ATN ✓ 70% decline in serum creatinine within 48 hrs of transplant ✓ Need of dialysis within 7 day of post transplant

48 Hrs

Sample collection and ELISA experiment

Five ml urine sample was collected serially at 0, 6,12,18,24 and 48 hrs of post transplant

Sample was centrifuged and supernatant was stored at -40 C

ELISA was performed with urine sample with R&D duo set ELISA kit.

Statistical Analysis

- Non parametric continuous variables were analyzed by using Mann Whitney U test.
- Categorical Variables was analyzed by Chie square test.
- Correlation was calculated by Pearson correlation coefficient ,two tailed test.
- ❖ Area under curve and receiver operating curve was plotted to determine the specificity, sensitivity and cutoff value of u Kim-1 for predicting DGF and ATN
- ❖ P value <0.05 was considered to be significant.

Results

Table 1 Demographic and clinical Profiles of Patient.

	-		%	
Characteristic		DGF(Mean±SD)	IGF(Mean±SD)	P -value
No. of patient		9	47	
Patient age(years)		38.0±12.94	39.68±11.86	0.702
Patient Weight(kg)		63.11±7.76	54.31±8.40	0.005
Patient Height(cm)		166.33±2.73	165.63±6.33	0.749
Patient BMI(kg/m²)		22.93±2.81	19.74±2.85	0.003
Patient.eGFR(mL/min/1.7	3m ²)at1week	40.35±14.43	65.39±16.91	<0.001
Donor's Age		44.00±18.17	47.87±9.29	0.481
Donor's GFR(mL/min/1.7	3 m ²)	41.89±12.05	45.10±14.84	0.546
Duration of dialysis (mont	hs)	10.77 ±4.11	11.36 ±4.32	0.710
Average seven days TLC		8.88 ±1.06	8.22 ±1.97	0.33
Average seven days serum		2.23 ±0.798	1.37 ±0.44	<0.001
creatinine(mg/dl)				
Average seven days Urine(liter)		5.21 ±1.98	7.29 ±2.19	0.011
Baseline creatinine		1.64 ±0.89	1.11 ±0.15	<0.001
Average seven days serum Tac		10.85±3.38	11.13±3.57	0.334
level(ng/ml)				
Cold ischemia time(minutes)		28.11±5.68	30.02±5.71	0.362
Warm ischemia time(minutes)		4.55±1.01	3.87±1.52	0.205
HLA mismatch score 0		0	6	0.499
	1	2	4	
2		1	10	
	3	3	17	

Tabel-2. Normalized uKIM-1/ creatinine ratio in pg/mg at different post transplant hours

Post transplant hours	DGF (Mean+SD)	IGF(Mean+SD)	P value		
0 Hrs	214.48±353.66	91.99±237.09	0.197		
6 Hrs	1521.88±639.53	623.76±474.90	<0.001		
12 Hrs	2201.39±810.096	934.91±831.74	<0.001		
18 Hrs	2863.15±809.11	792.72±296.21	<0.001		
24 Hrs	3258.76±912.57	796.93±186.68	<0.001		
48 Hrs	3967.45±850.04	761.38±221.52	<0.001		

Results

Table 3 Receiver operating area under curve (AUC-ROC) to

predict the delayed graft function and ATN.						
Time after	Area Under	Specificity	Sensitivity%	Predictive cut		
transplant	curve	%		off value		
				(pg/mg)		
0 Hrs	0.583	93.6	22.2	573.60		
6 Hrs	0.896	80.9	88.9	1031.19		
12 Hrs	0.896	83	88.9	1263.88		
18 Hrs	0.998	100	89.9	2012.36		
24 Hrs	1	100	89.9	2244.64		
表示是《····································	1 8	10000000	USSESSE TO	100 C		

Figure 2 AUC-ROC to predict the delayed graft function and ATN.

89.9

100

2871.42




Table 4. Urine volume at different Post transplant hours (ml)

Post transplant hours	DGF (Mean+SD)	IGF (Mean+SD)	P value	
>30 min	387.02±176.96	476.50±133.35	0.227	
6 Hrs	2167.33±322.59	2843.7±220.97	<0.001	
12 Hrs	4395.3±383.46	5420.30±287.51	< 0.001	
18 Hrs	7283.55±633.11	8470.0±518.65	< 0.001	
24 Hrs	9584.44±728.47	11112.20±1163.4	0.004	
48 Hrs	8035.22±1204.9	9845.6±932.28	0.002	

Table 5 Correlation of normalized uKIM-1(ng/mg) with confounding factors of donors

į,						242	211 223	
	Characteri	Patient.	Patient.	Patient.	Patient.	Seventh	Cold	Warm
	stics	Weight(Height	Age(Yea	BMI(Kg	day eGFR	ischemia	ischemia
		Kg)	(Cm)	rs)	/m ²)		time	time
	6 hours	r= 0.269	r=0.149	r=0.003	r=0.175	r=-0.415	r=0.018	r=0.174
	uKIM-1	p=0.05	p = 0.272	p=0.986	p=0.197	p=0.001	p=0.896	p=0.199
	12Hours	r=0.115	r=0.008	r=-0.011	r=0.096	r=-0.218	r=-0.099	r=0.258
	uKIM-1	p=0.40	p=0.951	p=0.933	p=0.480	p=0.106	p=.466	p=0.055
100	18 Hours	r=0.318	r=0.044	r=-0.009	r=0.327	r=-0.498	r=.105	r=0.176
300	uKIM-1	p=0.017	p=0.746	p=0.947	p=0.014	p<0.001	p=.440	p=0.194
	24Hours	r=0.386	r=0.046	r=0.012	r=0.397	r=-0.445	r=-0.020	r=0.117
	uKIM-1	p=0.003	p=0.736	p=0.932	p=0.002	p=001	p=.833	p=0.390
20	48Hours	r = 0.388	r=-0.093	r=-0.045	r=0.377	r=473	r=-0.074	r=0.223
	uKIM-1	p=0.003	p=0.494	p=0.743	p=0.004	p<0.001	p=0.589	p=0.098

Conclusions

- ✓ Urinary Kidney injury molecule -1 is a non invasive markers of delayed graft function and acute tubular necrosis.
- ✓ At 18 hours of post transplant, uKIM-1 level can predict DGF with 100% specificity and 88.9% sensitivity at cutoff value 2012.36pg/mg.
- ✓ The level of uKIM-1 excretion do not depend on demographic and other confounding factor of patient.
- ✓ It is highly specific and sensitive markers (within 18hrs) and can be used as marker for management of renal transplant patient patient with DGF and ATN.





ePosters

supported by

Roche Ltd.

F. Hoffmann-

