

The Star Excursion Balance Test as a measure of lower limb function in people with severe haemophilia

Sayers, Fionnuala., Cochrane, Judith., Benson, Gary. Haemophilia Comprehensive Care Centre, Belfast City Hospital, Belfast Health & Social Care Trust

Introduction and objectives

Current recommendations from the World Federation of Haemophilia recommend regular musculoskeletal assessment, incorporating Haemophilia Joint Health Score (HJHS) and Haemophilia Activities list (HAL) (Srivastava et al, 2013). However these tools lack appropriate measures of dynamic function to identify early functional deficits.

Common balance tests available include Berg Balance (Berg et al, 1992), timed "get-up and go" (Mathias et al, 1986), Functional reach (Duncan et al, 1990) and One-Legged-Stance (Fregly & Graybiel, 1968). These tests have limitations for those with early joint changes. Instrumented balance evaluation systems are expensive and time consuming.

The Star Excursion Balance Test (SEBT) has been shown to be a reliable non-instrumented test of lower extremity dynamic balance for some conditions (Gribble et al 2012). It involves balancing on one leg while reaching with the non-weight bearing leg in 3 directions (anterior, posterolateral and postero-medial). The aim of this study was to consider the suitability of the SEBT as a screening tool for patients with severe haemophilia.

Methods

10 patients with severe haemophilia A (aged 16–40 years) completed the SEBT as part of their annual musculoskeletal assessment. Reach distances were normalised to leg length and compared with ankle range of motion, HAL and HJHS. Pearson's test was used to identify relationships between reach distances, HAL, HJHS and range of movement.

References

Using the Star Excursion Balance Test to assess dynamic postural-control deficits and outcomes in lower extremity injury: A literature and systematic review. Gribble PA, Hertel J, Plisky P, Journal of Athletic Training (2012), 47(3):339-357.

Guidelines for the Management of hemophilia. Srivastava A, Brewer AK, Mauser-Bunschoten EP, Keys NS, Kitchen S, Llinas A, Ludlam CA, Mahlangu JN, Mulder K, Poon MC & Street A; Treatment guidelines working group on behalf of the World Federation of Hemophilia. Haemophilia (2013), 19, e1-e47

Measuring balance in the elderly: validation of an instrument. Berg KO, Wood-Dauphinee SL, Williams JJ, Maki B. Can.J.Public Health. 1992;83(Suppl 2):S7-11.

Balance in elderly patients: the "Get-up and go" test. Mathias S, Nayak US, Isaacs B. Arch.Phys.Med.Rehabil. 1986;67:387-389.

An ataxia test battery not requiring rails. Fregly AR, Graybiel A. Aerospace Medicine. 1968;39:277-282.

Functional reach: a new clinical measure of balance. Duncan PW, Weiner DK, Chandler J, Studenski S. Journal of Gerontology. 1990;45:M192-M197.

Results

20 Ankles

- Group A (n=9) normal ankles (no prior history of injury or bleed)
- Group B (n=11) target ankle joints


Group B demonstrated

- Reductions in mean reach distances in all directions for the SEBT compared with group A
- Reduction in mean ankle range of motion
- Poorer HJHS and HAL scores when compared with group A

For group B there was a strong positive correlation between HAL and anterior reach ($r^2=0.5732$, $p<0.01$), a moderate negative correlation between HJHS and anterior reach ($r^2=0.4779$, $p<0.01$), and a moderate positive correlation between dorsiflexion and anterior reach ($r^2=0.3602$, $p<0.01$). No relationship was demonstrated between HAL, HJHS or range of movement and posterolateral or posteromedial reach directions.

Conclusions

This study supports earlier work demonstrating that ankle dorsiflexion positively relates to performance of the SEBT. It also demonstrated a relationship between anterior reach distance and HJHS and HAL. It is quick and easy to administer requiring no more than a tape measure and approximately 2m² of floor space. As the test requires balance to be maintained at the limit of stability it may provide valuable information in lower extremity function in people with severe haemophilia.



Performance of the Star Excursion Balance test using the right limb as the stance limb in the (A) anterior, (B) posterolateral and (C) posteromedial directions.

